



## RESEARCH ARTICLE

10.1002/2014GC005658

## A global model for cave ventilation and seasonal bias in speleothem paleoclimate records

Eric W. James<sup>1</sup>, Jay L. Banner<sup>1</sup>, and Benjamin Hardt<sup>1,2</sup><sup>1</sup>Department of Geological Sciences and Environmental Science Institute, University of Texas at Austin, Austin, Texas, USA,<sup>2</sup>Now at Department of Earth, Atmospheric, and Planetary Sciences, Massachusetts Institute of Technology, Cambridge, Massachusetts, USA

## Key Points:

- Seasonal CO<sub>2</sub> variations in caves cause variations in speleothem growth rates. This can bias speleothem environmental proxies
- Seasonal ventilation should be strongest in mid to high latitudes relative to low latitudes, and in continental relative to maritime settings

## Supporting Information:

- Supporting Information S1

## Correspondence to:

E. W. James,  
ewjames@esi.utexas.edu

## Citation:

James, E. W., J. L. Banner, and B. Hardt (2015), A global model for cave ventilation and seasonal bias in speleothem paleoclimate records, *Geochem. Geophys. Geosyst.*, 16, 1044–1051, doi:10.1002/2014GC005658.

Received 20 NOV 2014

Accepted 6 MAR 2015

Accepted article online 16 MAR 2015

Published online 4 APR 2015

**Abstract** Cave calcite deposits (speleothems) provide long and continuous records of paleoenvironmental conditions in terrestrial settings. Typical environmental proxy measurements include speleothem growth rate and variations in elemental and isotope geochemistry. Commonly the assumption is made that speleothems grow continuously and at a constant rate throughout the year. However, seasonal variation of growth rate may be the rule in many caves. Here we apply observations of modern calcite growth and cave-air CO<sub>2</sub> concentrations and a model of factors controlling cave ventilation to construct a global model predicting where cave calcite growth may be seasonal. Previous models and measurements of calcite precipitation in caves demonstrate the retardation of speleothem growth by high cave-air CO<sub>2</sub>. Elevated CO<sub>2</sub> is commonly dissipated by ventilation driven by density differences between cave and surface air. Seasonal cycles in atmospheric temperature, pressure, and humidity commonly drive these density contrasts. Modeling these changes latitudinally and globally indicates a geographic control on seasonal cave ventilation and thus on a principal controlling factor of speleothem growth. The model predicts that given constant water, calcium, and CO<sub>2</sub> inputs, speleothems from temperate to boreal continental regions commonly accumulate more calcite in the cool season and less or none in the warm season. These models predict that proxies from temperate to boreal speleothems may be seasonally biased due to seasonal ventilation, whereas tropical and maritime records should not.

## 1. Introduction

Speleothem climate proxies provide valuable insights into past climates. Interpretation of these proxy results often assumes consistent growth throughout the year, providing an accurate depiction of mean annual climate. Recent studies of cave environments, however, suggest that this assumption may not be valid for many cave settings [Banner *et al.*, 2007; Baldini, 2010; Cowan *et al.*, 2013]. A comprehensive evaluation of the impacts of temperature-controlled density-driven ventilation can help distinguish regions that might be more prone to seasonal growth bias and help improve interpretation of paleoclimate records.


## 1.1. Controls on Growth Rate

The important equilibrium controls on precipitation of calcite include temperature [Plummer and Busenberg, 1982] and the chemical species in the calcite precipitation reaction



Increasing Ca<sup>2+</sup> or decreasing CO<sub>2</sub> drives the reaction to the right, precipitating calcite. Increasing CO<sub>2</sub> or decreasing Ca<sup>2+</sup> drives the reaction to the left, inhibiting precipitation. The equilibrium diagram for CO<sub>2</sub> and Ca concentrations shows the CO<sub>2</sub> contents required to allow or prohibit calcite precipitation (Figure 1). Although CO<sub>2</sub> levels of 2–5% have been measured in many caves, lower concentrations are the norm [EK and Gewalt, 1985; Banner *et al.*, 2007; Baldini, 2010].

The kinetics of speleothem calcite growth are governed by the extent of calcite oversaturation, the thickness and fluid dynamics of the water film that is precipitating calcite, drip rate, a kinetic constant related to temperature, and cave-air CO<sub>2</sub> level [Buhmann and Dreybrodt, 1985; Dreybrodt, 1988; Dreybrodt *et al.*,




**Figure 1.** Saturation diagram for equation (1). Shaded area shows the typical range of Ca concentrations in central Texas cave waters. Corresponding CO<sub>2</sub> equilibrium concentrations are enclosed by the long-dashed lines at 1 and 5.5%.

ing trace elements (e.g., Mg). It is important to note that CO<sub>2</sub> is only part of the equation, as changes in the Ca concentration will change growth rate as well.

### 1.2. CO<sub>2</sub> Gains, Losses, and Accumulation

Given the importance of CO<sub>2</sub> in controlling speleothem growth, we consider the sources and sinks of cave-air CO<sub>2</sub>. Microbial respiration in the unsaturated zone below the soil provides a CO<sub>2</sub>-rich source of air in some settings [Atkinson, 1977; Wood and Petraitis, 1984; Matthey et al., 2013]. In other caves, CO<sub>2</sub> inputs have been attributed to plant and microbial respiration in overlying soils [Ek and Gewalt, 1985; Spötl et al., 2005; Baldini et al., 2008; Breecker et al., 2012]. Typically, soil CO<sub>2</sub> concentration correlates positively with soil temperature and water availability [Harper et al., 2005]. Soil CO<sub>2</sub> concentrations are highest in midsummer in climates with sufficient effective precipitation year-round, but peaks in the spring to early summer in regions where soils dry out during the summer. In drips fed directly from soil water, these high-CO<sub>2</sub> periods should be periods of the highest dissolution of carbonate bedrock and thus the highest delivery of Ca and HCO<sub>3</sub><sup>-</sup> (and ultimately CO<sub>2</sub>) to caves.




**Figure 2.** Calculated and measured growth rates of speleothems with respect to cave air CO<sub>2</sub> concentrations. Dashed, solid, and gray lines show calculated rates [Buhmann and Dreybrodt, 1985; Dreybrodt, 1988; Dreybrodt et al., 1997] at three Ca concentrations. Conditions are 20°C, one drop per second, water film thickness of 0.5 mm, a kinetic constant of  $2.6 \times 10^{-5}$  cm/s, and apparent Ca equilibrium factor of 1.195. Diamonds and circles indicate measured growth rates and CO<sub>2</sub> concentrations from two drip water sites in Inner Space Cavern (ISST and ISLM) where calcite growth on artificial substrates is monitored, Table S1.

1997; Baker et al., 2014]. Theoretical growth rates (Figure 2) are approximately halved when the ambient CO<sub>2</sub> level is increased from atmospheric (390 ppm) to 2500 ppm in water containing 2 mmol/L Ca. Comparisons of measured CO<sub>2</sub> concentration and calcite growth rate in cave environments are consistent with these models (Figure 2, Table S1) [Baker et al., 1998, 2014]. Note that cessation of calcite deposition occurs in several Texas caves when CO<sub>2</sub> concentrations are more than 5000 ppm [Banner et al., 2007]. Calculated growth rates that differ from measured growth rates may be due to incomplete capture of precipitating calcite [Banner et al., 2007], microbial mediation of calcite deposition [Barton and Northup, 2007], and uncertainties in the kinetic constant or in estimating water thickness [Baker et al., 2014] or interfering

trace elements (e.g., Mg). It is important to note that CO<sub>2</sub> is only part of the equation, as changes in the Ca concentration will change growth rate as well. Although soil and unsaturated zone CO<sub>2</sub> flux to caves is an important variable controlling Ca and CO<sub>2</sub> delivery, the concentration of Ca in drip waters does not necessarily follow the seasonal cycling of CO<sub>2</sub> [Tooth and Fairchild, 2003; Banner et al., 2007]. Calcium concentration and water availability are commonly influenced by independent variables such as water storage and transit time.

If caves are closed to the atmosphere, then CO<sub>2</sub> should accumulate until calcite precipitation stops [Moore and Nicholas, 1964, p. 34]. Few caves, if any, are gas-tight systems. We emphasize the importance of seasonal cave ventilation, which is commonly a consequence of density differences between air inside and



**Figure 3.** Modeled seasonal air densities in caves and at the surface for three locations: Minnesota, Texas, and Guam. Daily minimum (blue line) and maximum (red) surface air densities compared to cave air density estimated from average yearly temperature, daily barometric pressure, and 95% RH (black) for each location [NOAA, 2010]. When the maximum daily surface air density (red) is below the cave air density (black), there is no density-driven exchange of air between cave and surface. This period of potential  $\text{CO}_2$  accumulation is highlighted in yellow in Figure 3a. When surface air density is above cave air density, there is density-driven flow of outside air into the cave, flushing the cave with  $\sim 390$  ppm  $\text{CO}_2$  air. (c) In tropical settings the diurnal density range (red to blue lines) brackets cave air density, suggesting diurnal ventilation that would prevent  $\text{CO}_2$  accumulation due to air stagnation. Note that the duration and magnitude of summer density differences are greatest at (a) the highest latitude site. Purple line in Figure 3b shows seasonal changes in calcite precipitation rate for site ISST, Table S1.

[Moore and Nicholas, 1964]. Additionally, RH in the deep parts of most caves, particularly those containing growing speleothems, is high and does not vary substantially with surface RH [Wigley and Brown, 1976]. It can be assumed that cave air pressure typically adjusts to surface pressure within a matter of hours. Surface T, P, and RH cycle seasonally, synoptically, and diurnally. The combination of relatively constant cave temperature approximating the local annual mean with the variation of local surface temperatures above and below this mean commonly results in predictable seasonal airflow. Thus, differences in T, P, RH, and composition between cave air and surface air create differences in density that can (1) stratify cave and surface air (2) mix cave and surface air, or (3) force less dense air out of a cave through the primary entrance, secondary entrances, or fracture systems.

### 1.3. The Effects of Seasonal Temperature Cycles

As the year progresses into fall, daily minimum surface temperatures decrease and drop below the temperature of the cave air. At this time, air density outside the cave becomes greater than that inside the cave. The denser, cooler outside air flows into the cave as the warmer, lighter cave air flows out. High  $\text{CO}_2$  air is flushed out of the cave and replaced by low  $\text{CO}_2$  air ( $\sim 390$  ppm). The comparatively low  $\text{CO}_2$  content of the outside air allows faster calcite precipitation. At some time in the spring, daily low temperatures rise above the cave air temperature and cave and surface air stratify due to density differences.  $\text{CO}_2$  concentrations in the cave will begin to build and rise through the summer, bolstered by increasing supply from the warming soil.

This yearly cycle of cave-air  $\text{CO}_2$  concentrations can be modeled for most caves using local weather data [NOAA, 2010], T, P, and RH, to calculate outside air densities (Figure 3). The density of cave air can be calculated if T, P, and RH have been measured or can be approximated with annual average temperature, an approximated RH (80–100%), and measured outside atmospheric pressure. At mid to high latitudes this yields a surface air density versus time record with diurnal density variations superimposed on a yearly surface-air density cycle. This yearly cycle has a density minimum in the summer and a maximum in winter

outside a cave, as one important cause of seasonal cycling of cave  $\text{CO}_2$  concentrations [Wigley and Brown, 1976; Buecher, 1999]. Air density is a function of temperature (T), pressure (P), relative humidity (RH), and gas composition. For nontropical latitudes, temperature is the most important of these variables over a seasonal cycle. A change from 25 to 0°C at 65% humidity results in approximately an 11% increase in air density. A 3 kPa increase in P at 20°C, by contrast, yields only a 3.2% increase in density. A change from 100% RH to 0% at 20°C increases density by less than 1%. An increase in the  $\text{CO}_2$  concentration from 400 ppm to 1% at 20°C and 100% RH increases air density by just over 0.5% [Sánchez-Cañete et al., 2013]. Outside T and RH are progressively damped as one moves into a cave [Wigley and Brown, 1976] and at sufficient depth cave temperature represents the local average yearly temperature

(Figures 3a and 3b). In comparison, cave air density is relatively constant throughout the year. The yearly surface air density cycle crosses the cave air density curve in the spring and the fall. The period in the summer when cave air density is greater than surface air density typically becomes longer and the density difference more profound with increasing latitude of the site (Figure 3).

When the surface air density is less than the cave air density, caves with chambers below the entrance should experience CO<sub>2</sub> buildup during these periods. This seasonal cyclicity in CO<sub>2</sub> has been observed in caves in many regions, including southwestern USA [Buecher, 1999; Banner *et al.*, 2007; Cowan *et al.*, 2013] (Figure 3a), southeastern USA [Kowalczyk and Froelich, 2010], and Europe [Spötl *et al.*, 2005; Bourges *et al.*, 2006; Liñán *et al.*, 2008; Baldini, 2010]. This summer CO<sub>2</sub> buildup (Figure 3) and the inhibition of calcite precipitation by elevated CO<sub>2</sub> levels (Figure 2) can cause a seasonal bias in speleothem growth. There will be more cool-season calcite precipitation than warm-season calcite precipitation [Banner *et al.*, 2007; Baldini *et al.*, 2008; Baldini, 2010]. Additionally, speleothem growth rate changes, caused by ventilation, can cause seasonal differences in C and O isotope compositions of calcite [Spotl *et al.*, 2005; Scholz *et al.*, 2009; Baldini, 2010; Feng *et al.*, 2012]. Because seasonal growth in most speleothems commonly eludes observation or sampling resolution, such biases can be significant. Proxy measurements from speleothems that grew in caves where summer CO<sub>2</sub> is elevated will yield measurements biased toward winter conditions in proportion to winter versus summer calcite growth. However, if these seasonal variations can be observed they become a chronologic tool.

#### 1.4. Exceptions to Warm Season CO<sub>2</sub> Accumulation


We expect that this type of cyclicity in CO<sub>2</sub> concentration is common, but certainly not universal. If the cave entrance is at a lower elevation than the cave itself, summer and winter ventilation regimes will be reversed. The large class of caves with multiple openings has complex ventilation patterns [e.g., Bourges *et al.*, 2006; Spötl *et al.*, 2005]. Those with openings at different elevations may change ventilation direction seasonally due to a “chimney effect,” whereby air enters and exits at different openings [Wigley and Brown, 1976; Atkinson *et al.*, 1983]. In some settings, seasonal winds may cause summer ventilation. This is governed by wind direction and the orientation of the cave and its entrances [Kowalczyk and Froelich, 2010]. Flowing water in caves can influence local T, RH, and pCO<sub>2</sub> significantly. CO<sub>2</sub> ventilation is less efficient in caves with tortuous or narrow passages and in very deep caves [Moore and Nicholas, 1964; Wigley and Brown, 1976]. Low effective moisture in summer may limit CO<sub>2</sub> production and transportation, and climates with strong winter vegetation growth may supply more CO<sub>2</sub> to caves in winter. In the tropics, the diurnal temperature range, and thus density difference, is greater than seasonal density fluctuations and therefore the temperature-driven ventilation model does not predict a seasonal cycle (Figure 3c).

It should be noted that our ventilation model only addresses the seasonal output of CO<sub>2</sub> from caves; it does not take into account seasonal variations in CO<sub>2</sub> supply or the positive feedback between increasing CO<sub>2</sub> concentration and air density [Sánchez-Cañete *et al.*, 2013]. However, both of these influences should act to amplify CO<sub>2</sub> accumulation in a seasonal time frame. CO<sub>2</sub> supply typically reflects soil productivity and is thus highest in spring and summer. Increasing cave CO<sub>2</sub> acts to increase cave air density, thus suppressing ventilation further. In some parts of the tropics, strong seasonality in rainfall and weak seasonality in temperature may result in soil CO<sub>2</sub> production producing seasonal cave-air CO<sub>2</sub> fluctuations, whereas our ventilation model does not predict a seasonal cycle.

## 2. Methods: Geographic Variation in Seasonal Cave Ventilation

To assess geographic variations in seasonal cave ventilation at the global scale, we construct a map (Figure 4) of the average yearly sum of the positive differences between average monthly minimum temperature and the annual mean temperature. These temperature differences ( $\Delta T_m$ ) are essentially the summer temperature contrast between the surface and a cave, which accounts for virtually all the density differences, as depicted in yellow highlight in Figure 3a.

$\Delta T_m$  is constructed using the sites in the Global Historical Climatology Network (GHCN) monthly data set v. 3 [NOAA, 2011], for which there were complete years (12 months) of temperature values from 1980 to 2010. An annual station mean was calculated for each year in the data set from 1980 to 2010 by taking the average of the monthly means weighted for the number of days in the month. The annual mean temperature



**Figure 4.** Map of temperature differences,  $\Delta T_m$ , which represent the temperature contribution to air density contrasts (Figure 3a, yellow highlight), which typically drive cave ventilation. Because temperature is the major seasonally cyclic control of air density this is a first-order map of the seasonal differences between cave air densities and surface air densities. Cave atmospheres in regions with higher  $\Delta T_m$  are more likely to stagnate seasonally and accumulate  $\text{CO}_2$ , which reduces calcite growth rates.  $\Delta T_m$  is calculated as the average yearly sum of the positive differences between average monthly minimum temperature and the annual mean temperature for 1980–2010. No elevation corrections have been applied. Data are from GHCN [NOAA, 2011]. Data from Poland prior to 1995 are anomalously low and have been excluded.

for that year was subtracted from the average monthly minimum temperatures. These monthly differences were then averaged by station for all years. The annual number was determined by summing the differences for months when the average monthly minimum temperature exceeded the annual mean, giving the value  $\Delta T_m$ . The units of  $\Delta T_m$  are  $^{\circ}\text{C}$  months.

### 3. Results

#### 3.1. Latitudinal Structure

$\Delta T_m$  is a good representation of the temperature contribution to the density differences between cave air and surface air in the summer months, as highlighted in yellow in Figure 3a. Thus, Figure 4 provides a first-order approximation of the strength and duration of seasonal stagnation of cave air. This indicates regions where ventilation-controlled  $\text{CO}_2$  accumulation in caves is potentially important in controlling speleothem growth. In areas where  $\Delta T_m$  values are 0–3 $^{\circ}\text{C}$  months (Figure 4, red symbols) the diurnal temperature range typically overlaps the annual temperature variation. This drives diurnal density changes in surface air that range above and below cave air density, suggesting daily ventilation, and thereby allowing year-round calcite growth. Moving poleward or toward continental interiors to orange to yellow to green ( $\Delta T_m = 3$ –12  $\Delta T_m$ ), blues (12–15  $\Delta T_m$ ), and indigo regions ( $>15$   $\Delta T_m$ ), there is an increasing likelihood that the hottest periods of the year will result in minimum surface temperatures that are above the mean annual temperature, which approximates local cave temperature. Without large surface versus subsurface differences in pressure or humidity affecting air densities, there will be little or no summer cave ventilation in these regions.

#### 3.2. Local Geographic Structure

Figure 4 shows several significant deviations from a simple latitudinal trend. The first is where maritime climatic influences moderate annual temperature extremes. Coastal sites commonly have yearly air density curves that allow density-driven ventilation on shorter time spans than the seasons. For example, the temperature range above and below the yearly average is similar along the North American west coast and in

the tropics to subtropics. Thus, in tropical and maritime settings CO<sub>2</sub> concentrations in caves are likely controlled by variations in supply from soil, diurnal temperature variation, synoptic pressure changes, and winds. CO<sub>2</sub> levels in soils commonly peak in summer in mid to high latitudes, paralleling ventilation-controlled cave CO<sub>2</sub> buildup. Other deviations from latitudinal control are in continental deserts and at high elevations, which typically have large diurnal temperature ranges that would drive diurnal ventilation and therefore limit seasonal effects.

## 4. Discussion

### 4.1. Geographic Variations in Ventilation

The exchange of cave air with surface air, along with variations in CO<sub>2</sub> input and water supply, are important controls on cave-air CO<sub>2</sub> and the therefore the growth of speleothem calcite. The principal driving force in cave ventilation is the density contrast between cave air and surface air, and this contrast is a function of air temperature, pressure, and relative humidity. Winter cave ventilation and summer cave-air stagnation are consistent with model results and observations from midlatitude sites of faster speleothem calcite growth in winter than in summer. These results yield a global model that predicts the potential for air exchange, cave-air CO<sub>2</sub> variation, and seasonal speleothem growth. The model predicts that caves in the tropics ventilate diurnally. Similarly, maritime climates dampen seasonal temperature extremes and, therefore, seasonal ventilation effects. Changes in CO<sub>2</sub> supply and subseasonal ventilation due to synoptic and subsynoptic scale weather are the likely controls of CO<sub>2</sub> levels in caves in tropical and maritime settings. In contrast, caves in temperate to boreal regions and in continental interiors are predicted to preferentially ventilate in the cool season.

### 4.2. Implications for Speleothem Paleoclimate Records

The global model predicts a lack of summer cave ventilation and consequent accumulation of CO<sub>2</sub> in caves at mid to high latitudes, which results in a summer slowing or cessation of speleothem growth. This prediction implies that paleoclimate reconstructions from speleothems from temperate to boreal regions and continental interiors are more likely to be seasonally biased toward recording cool season conditions, whereas those from tropical and maritime regions likely grow more continuously. The model yields a map (Figure 4) that can be used to predict this potential bias toward preservation of cool season conditions in speleothem proxies as a consequence of differences in density-driven cave ventilation. Comparisons of proxy records between caves in regions with different climates may be particularly problematic. Furthermore, even on a local scale the seasonality of ventilation may affect the geochemical parameters typically used as paleoclimate proxies at some drip sites more than at others [Wong *et al.*, 2011]. In view of the exceptions to the general geographic pattern [e.g., Spötl *et al.*, 2005; Matthey *et al.*, 2008] (also see section 1.4) the seasonal airflow should be considered on a site-by-site basis.

### 4.3. Capture of Seasonal Information

Seasonal growth bias has both negative and positive consequences for speleothem climate proxies. An example of the potential for missing climatic data is illustrated by likely differences in the record of storm events in speleothems of tropical versus subtropical locations. Frappier *et al.* [2007] document the passing of individual tropical cyclones using the oxygen isotope records in fast-growing speleothems from Belize (Figure 4). However, data from central Texas caves (Figure 4) indicate that there is high CO<sub>2</sub> and little or no growth [Banner *et al.*, 2007] in central Texas during the hurricane season. For example, in Inner Space Cavern, only 2% of the annual growth takes place from August through October [Banner *et al.*, 2007] when 95% of major Atlantic hurricanes occur [Landsea, 1993]. The applicability of this paleotempestological technique is restricted to caves that allow the growth of summer and early fall calcite or in rare instances when large storms deliver a significant fraction of annual precipitation.

Seasonal patterns of growth are not exclusively an impediment to developing environmental proxy records. When recognized and understood, they are signal rather than noise. Speleothem calcite texture, trace element content, and stable isotopes values are controlled in part by cave air CO<sub>2</sub> concentration [Stoll *et al.*, 2012; Matthey *et al.*, 2008, 2010; Feng *et al.*, 2012]. Additionally, seasonal growth can provide a basis for layer-counting to build multiyear chronologies and as a means of focusing on an interval within a year, increasing the proxy's resolution, albeit at the expense of knowledge about periods of less growth. The timing and

duration of seasonal speleothem growth is a proxy for the intensity and duration of seasons (Figure 3). Measuring such differences is typically near the limit of analytical capabilities, but improved methods coupled with samples with fast growth rates may allow quantitative measures of seasonality.

## 5. Conclusions

Global-scale climate drives ventilation in many caves, and this seasonal ventilation is one control of speleothem growth rate. Seasonal ventilation should be strongest in mid to high latitudes relative to low latitudes, and in continental relative to maritime settings. In most caves, seasonal growth bias is toward the preservation of cooler season records. Furthermore, seasonal growth patterns in temperate to boreal latitudes may preclude the identification of events such as hurricanes that occur during a specific time of year. This model of cave air ventilation emphasizes the importance of understanding a cave system's ventilation dynamics prior to interpretation of paleoclimate records.

### Acknowledgments

This work was supported by the US National Science Foundation grants ATM-0823665 to Banner and Yang, EAR-1124514 to Breecker and Banner and by the Geology Foundation and the Environmental Science Institute of The University of Texas. Data sources are Table S1 for Figure 2, Table S1 and online data [NOAA, 2010] for Figure 3 and online data [NOAA, 2011] for Figure 4. We thank T. Atkinson and an anonymous reviewer for their constructive comments.

### References

- Atkinson, T. C. (1977), Carbon dioxide in the atmosphere of the unsaturated zone: An important control of groundwater hardness in limestones, *J. Hydrol.*, *35*, 111–123.
- Atkinson, T. C., P. L. Smart, and T. M. L. Wigley (1983), Climate and natural radon levels in Castleguard cave, Columbia icefields, Alberta, Canada, *Arct. Alp. Res.*, *15*, 487–502.
- Baker, A., D. Genty, W. Dreybrodt, W. L. Barnes, N. J. Mockler, and J. Grapes (1998), Testing theoretically predicted stalagmite growth rate with recent annually laminated samples: Implications for past stalagmite deposition, *Geochim. Cosmochim. Acta*, *62*, 393–404, doi:10.1016/S0016-7037(97)00343-8.
- Baker, A. J., Matthey, D. P., and J. U. L. Baldini (2014), Reconstructing modern stalagmite growth from cave monitoring, local meteorology, and experimental measurements of dripwater films, *Earth Planet. Sci. Lett.*, *392*, 239–249.
- Baldini, J. U. L. (2010), Cave atmosphere controls on stalagmite growth rate and paleoclimate records, in *Tufas and Speleothems: Unraveling the Microbial and Physical Controls*, *Spec. Publ. 336*, edited by H. Perley and M. Rogerson, pp. 283–394, Geol. Soc., London, U. K., doi:10.1144/SP336.15.
- Baldini, J. U. L., F. McDermott, D. L. Hoffmann, D. A. Richards, and N. Clipson (2008), Very high-frequency and seasonal cave atmosphere PCO<sub>2</sub> variability: Implications for stalagmite growth and oxygen isotope-based paleoclimate records, *Earth Planet. Sci. Lett.*, *272*, 118–129, doi:10.1016/j.epsl.2008.04.031.
- Banner, J. L., A. Guilfoyle, E. W. James, L. A. Stern, and M. Musgrove (2007), Seasonal variations in modern speleothem calcite growth in central Texas, U.S.A., *J. Sediment. Res.*, *77*, 615–622, doi:10.2110/jsr.2007.065.
- Barton, H. A., and D. E. Northup (2007), Geomicrobiology in cave environments: Past, current and future perspectives, *J. Cave Karst Stud.*, *69*, 163–178.
- Bourges, F., P. Genthon, A. Mangin, and D. d'Hulst (2006), Microclimates of l'Aven d'Orgnac and other French limestone caves (Chauvet, Esparros, Marsoulas), *Int. J. Climatol.*, *26*, 1651–1670, doi:10.1002/joc.1327.
- Breecker, D. O., A. E. Payne, J. Quade, J. L. Banner, C. E. Ball, K. W. Meyer, and B. D. Cowan (2012), The sources and sinks of CO<sub>2</sub> in caves under mixed woodland and grassland vegetation, *Geochim. Cosmochim. Acta*, *96*, 230–246, doi:10.1016/j.gca.2012.08.023.
- Buecher, R. (1999), Microclimate study of Kartchner Caverns, Arizona, *J. Cave Karst Stud.*, *61*, 108–120.
- Buhmann, D., and W. Dreybrodt (1985), The kinetics of calcite dissolution and precipitation in geologically relevant situations of karst areas: 1. Open system, *Chem. Geol.*, *48*, 189–211, doi:10.1016/0009-2541(85)90046-4.
- Cowan, B., M. Osborne, and J. Banner (2013), Temporal variability of cave-air CO<sub>2</sub> in central Texas, *J. Cave Karst Stud.*, *75*, 38–50, doi:10.4311/2011ES0246.
- Dreybrodt, W. (1988), *Processes in Karst Systems: Physics, Chemistry, and Geology*, Springer, Berlin.
- Dreybrodt, W., L. Eisenlohr, B. Madry, and S. Ringer (1997), Precipitation kinetics of calcite in the system CaCO<sub>3</sub> - H<sub>2</sub>O - CO<sub>2</sub>: The conversion to CO<sub>2</sub> by the slow process H<sup>+</sup> + HCO<sub>3</sub><sup>-</sup> - CO<sub>2</sub> + H<sub>2</sub>O as a rate limiting step, *Geochim. Cosmochim. Acta*, *61*, 3897–3904, doi:10.1016/S0016-7037(97)00201-9.
- Ek, C., and M. Gewalt (1985), Carbon dioxide in cave atmospheres. New results in Belgium and comparison with some other countries, *Earth Surf. Processes Landforms*, *10*, 173–187, doi:10.1002/esp.3290100209.
- Feng, W., J. L. Banner, A. L. Guilfoyle, M. Musgrove, and E. W. James (2012), Oxygen isotopic fractionation between drip water and speleothem calcite: A 10-year monitoring study, *Chem. Geol.*, *304*(305), 53–67, central Texas, USA.
- Frappier, A., D. Sahagian, S. Carpenter, L. Gonzalez, and B. Frappier (2007), Stalagmite stable isotope record of recent tropical cyclone events, *Geology*, *35*, 111–114.
- Harper, C. W., J. M. Blair, P. A. Fay, A. K. Knapp, and J. D. Carlisle (2005), Increased rainfall variability and reduced rainfall amount decreases soil CO<sub>2</sub> flux in a grassland ecosystem, *Global Change Biol.*, *11*, 322–334, doi:10.1111/j.1365-2486.2005.00899.x.
- Kowalczyk, A. J., and P. N. Froelich (2010), Cave air ventilation and CO<sub>2</sub> outgassing by radon-222 modeling: How fast do caves breathe?, *Earth Planet. Sci. Lett.*, *289*, 209–219.
- Landsea, C. W. (1993), A climatology of intense (or major) Atlantic hurricanes, *Mon. Weather Rev.*, *121*, 1703–1713, doi:10.1175/1520-0493(1993)121<1703:ACOIMA>2.0.CO;2.
- Liñán, D., I. Vadillo, and F. Carrasco (2008), Carbon dioxide concentration in air within the Nerja Cave (Malaga, Andalusia, Spain), *Int. J. Speleol.*, *37*, 99–106.
- Matthey, D., D. Lowry, J. Duffet, R. Fisher, E. Hodge, and S. Frisia (2008), A 53 year seasonally resolved oxygen and carbon isotope record from a modern Gibraltar speleothem: Reconstructed drip water and relationship to local precipitation, *Earth Planet. Sci. Lett.*, *269*, 80–95, doi:10.1016/j.epsl.2008.01.051.
- Matthey, D. P., I. J. Fairchild, T. C. Atkinson, J. P. Latin, M. Ainsworth, and R. Durell (2010), Seasonal microclimate control of calcite fabrics, stable isotopes and trace elements in modern speleothem from St Michaels Cave, Gibraltar, *Geol. Soc. Spec. Publ.*, *336*, 323–344, doi:10.1144/SP336.17.

- Mattey, D. P., R. Fisher, T. C. Atkinson, J. P. Latin, R. Durrell, M. Ainsworth, D. Lowry, and I. J. Fairchild (2013), Methane in underground air in Gibraltar karst, *Earth Planet. Sci. Lett.*, *374*, 71–80.
- Moore, G., and G. Nicholas (1964), *Speleology: The Study of Caves*, Heath and Co., Boston.
- NOAA (2010), National Climate Data Center Data Online. [Available at <http://cdo.ncdc.noaa.gov/pls/plclimprod/cdomain.abbrev2id>.]
- NOAA (2011), Global Historical Climatology Network—Monthly Version 3. [Available at <http://www.ncdc.noaa.gov/ghcnm/v3.php>.]
- Plummer, L. N., and E. Busenberg (1982), The solubilities of calcite, aragonite and vaterite in CO<sub>2</sub>-H<sub>2</sub>O solutions between 0 and 90°C, and an evaluation of the aqueous model for the system CaCO<sub>3</sub>-CO<sub>2</sub>-H<sub>2</sub>O, *Geochim. Cosmochim. Acta*, *46*, 1011–1040, doi:10.1016/0016-7037(82)90056-4.
- Sánchez-Cañete, E., P. Serrano-Ortiz, F. Domingo, and A. Kowalski (2013), Cave ventilation is influenced by variations in the CO<sub>2</sub>-dependent virtual temperature, *Int. J. Speleol.*, *42*, 1–8.
- Scholz, D., C. Mühlinghaus, and A. Mangini (2009), Modelling  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  in the solution layer on stalagmite surfaces, *Geochim. Cosmochim. Acta*, *73*, 2592–2602.
- Spötl, C., I. J. Fairchild, and A. F. Tooth (2005), Cave air control on dripwater geochemistry, Obir Caves (Austria): Implications for speleothem deposition in dynamically ventilated caves, *Geochim. Cosmochim. Acta*, *69*, 2451–2468, doi:10.1016/j.gca.2004.12.009.
- Stoll, H. M., W. Müller, and M. Prieto (2012), I-STAL, a model for interpretation of Mg/Ca, Sr/Ca and Ba/Ca variations in speleothems and its forward and inverse application on seasonal to millennial scales, *Geochim. Geophys. Geosyst.*, *13*, Q09004, doi:10.1029/2012GC004183.
- Tooth, A. F. and I. J. Fairchild (2003), Soil and karst aquifer hydrological controls on the geochemical evolution of speleothem-forming dripwaters, Crag Cave, southwest Ireland, *J. Hydrol.*, *273*, 51–68.
- Wigley, T. M. L., and M. C. Brown (1976), The physics of caves, in *The Science of Speleology*, vol. 3, edited by T. D. Ford and C. H. D. Cullingford, pp. 329–347, Academic, N. Y.
- Wong, C. I., J. L. Banner, and M. Musgrove (2011), Seasonal dripwater Mg/Ca and Sr/Ca variations driven by cave ventilation: Implications for and modeling of speleothem paleoclimate records, *Geochim. Cosmochim. Acta*, *75*, 3514–3529, doi:10.1016/j.gca.2011.03.025.
- Wood, W. W., and M. J. Petraitis (1984), Origin and distribution of carbon dioxide in the unsaturated zone of the southern high plains of Texas, *Water Resour. Res.*, *20*, 1193–1208.