

4 miles east of Diboll on Stovall Creek.  
(Loc.12 of Cushman and Applin, Bull. AAPG,  
vol.10, p.185, 1926)

Type locality for--

- × Textularia dibollensis Cushman & Applin (Bull.AAPG, vol.10, p.165, pl.6, figs.12-14, 1926)
- × Bolivina gracilis Cushman & Applin ( Idem, p.167, pl.7, figs.1,2)
- × Bolivina jacksonensis var. striatella Cushman & Applin (Idem, p.167, pl.7, figs.5,6)
- × Nodosaria (Glandulina) laevigata var. ovata Cushman & Applin (Idem, p.169, pl.7, figs.12,13)
- × Discorbis jacksonensis var. dibollensis Cushman & Applin (Idem, p.178, pl.9, fig.10)
- × Truncatulina americana var. antiqua Cushman & Applin (Idem, p.179, pl.9, figs.12,13)
- × Anomalina granosa var. dibollensis Cushman & Applin (Idem, p.179, pl.9, fig.15)
- × Siphonina jacksonensis Cushman & Applin (Idem, p.180, pl.9, figs.20-23)
- × Pulvinulina jacksonensis Cushman & Applin (Idem, p.181, pl.9, figs.24,25)
- × Nonionina advena var. inexcavata Cushman & Applin (Idem, p.182, pl.10, figs.18,19)

Exposure on White Oak Creek, just below  
bridge on Club road,  $\frac{1}{2}$  mile SE. of Diboll.

Type locality for--

- x Textularia mississippiensis var. rhomboidea Cushman  
and Ellisor (C. C., vol.7, p.51, pl.7, fig.2, 1931)