

Midway group  
Wills Point fm.

-Road cut close to Escondido Creek about 1½ miles from its confluence with Martinez Creek (Floresville quadrangle, Military Sheet 467-S-II & IV). This clay carries a generous scattering of upper Midway foraminifera.

(Sta.82, Univ.Texas Bull.2644, p.62, 1927)

15-T-2  
(Bexar County)

Midway group  
Wills Point fm.

-Alamo Brick Company clay pit southeast of San Antonio (San Antonio topographic sheet) on north side of the road leading southeast from the city and about one-quarter of a mile from the area broken up into city blocks; just beyond the brick yard the road forks, one branch leading to Gonzales, the other to Sutherland Springs. The compact clay from the bottom of this excellent exposure washes down to a small residue that carries charas, otoliths, small gastropods, and a large number of foraminifera of the upper Midway faunal unit. *Allomorphina trigona* is especially abundant in the clays of this outcrop, and *Chilostomelloides eocenica* is a rare form.

15-T-3  
(Bexar County)

(Sta.85, Univ.Texas Bull.2644, p.62, 1927)

Type locality for --

Vitrewebbina chapmani Plummer (Univ. Texas Bull. 2644, p. 128, pl.8, fig.2, 1927)

Albmorphina halli Jennings (Bull. Amer. Paleon., #78, p.34, pl.4, fig.5, 1936; new name for A. trigona of Plummer, 1927, not Reuss, 1850).

Type locality for --  
*Vitrewebbina chapmani* Plummer

Midway group  
Wills Point fm.

-Road cut close to an acute angle in the road 2 miles southwest of Zuehl (Floresville quadrangle,<sup>17</sup> Military Sheet, 467-S-II & IV). An abundance of foraminifera characterize these clays, and they place these strata definitely in the upper part of the Midway formation.

—Sta. 79, Univ. Texas Bull. 2644, p. 62, 1927)

15-T-5  
(Bexar County)

Midway group  
Wills Point fm.

15-T-6  
(Bexar County)

—In bank of Salitrillo Creek about 1 mile from its junction with Martinez Creek (Floresville quadrangle, Military Sheet 467-S-II & IV). A great abundance of the upper Midway forms characterize the clays of this outcrop.

(Sta.80, Univ.Texas Bull.2644, p.62, 1927)

Midway group  
Wills Point fm.

15-T-7  
(Bexar County)

—Tank between Escondido and Salitrillo creeks about 1½ miles west of station 80 and east of the northeast-southwest road (Floresville quadrangle, Military Sheet 467-S-II & IV). An abundance of rather unusually thin-shelled and emaciate foraminiferal tests of the upper Midway faunule mark this outcrop.

L (Sta.81, Univ.Texas Bull.2644, p.62, 1927)

Taylor group  
(Upper part)

152. Ditch on second-class road leading down Saltillo Creek  
Valley  $3\frac{1}{2}$  miles (air line) southeast of Converse, Bexar  
County, Tex. L. W. Stephenson.

(P.P. 206)

15-T-10  
(Bexar County)



Type locality for --

Dorothia stephensoni Cushman (C.C., Sp. Pub. #6, p. 28)

Type loc. for --  
Dorothia stephensoni Cushman (C.C., Sp. Pub. #6, p. 28, fig. 36)

Taylor group  
(Upper part)

158. Gully in west-facing slope of Medio Creek Valley below  
second-class road  $\frac{1}{4}$  mile north of Castroville road,  
Bexar County, Tex. L. W. Stephenson.

(P.P. 206)

15-T-11  
(Bexar County)

Type locality for —

Gaudryina quadrans Cushman (C. Spec. Pub. 6, p. 6)

Collected for —  
Gaudryina quadrans Cushman (C. Spec. P. 6, p. 6, 36)

Navarro group  
Corsicana fm

Bank on south side of old San Antonio-Castroville highway (immediately south of the new highway), 17.3 miles by road west of the Missouri Pacific Railway station in San Antonio, or 6 miles east of Castroville, or 2.8 miles by road east of the Bexar-Medina county line. Stephenson records the Taylor-Navarro contact in the exposure several feet below the hard Exogyra-Gryphaea capping layer. Sample 3 feet below this ledge has furnished holotypes for two species.

15-T-12  
(Bexar County)

Type locality for--

- x Ammobaculites texanus Cushman (C.C., vol.9, p.50, pl.5, fig.3, 1933)
- x Buliminella carseyae var. plana Cushman & Parker (C.C., vol.12, p.8, pl.2, fig.7, 1936)

Navarro group  
Corsicana fm.

46. 3 feet below base of *Exogyra-Gryphaea* bed, San Antonio  
road 6 miles east of Castroville, Bexar County, Tex.  
L. W. Stephenson. (P. P. 266)

15-T-12  
(Bexar County)



Type locality for --

- 5-3 x *Ammobaculites texanus* Cushman (C.C., vol.9, p.50)  
2-7 x *Buliminella carseyae* var. *plana* Cushman and  
Parker (C.C., vol.12, p.8)

1936

(Plummer Sta 23)

Type loc. for --  
*Ammobaculites texanus* Cushman (C.C. vol.9, p.50)  
*Buliminella carseyae*  
var. *plana* Cushman & Parker (C.C. vol.12, p.8)