

Taylor group
(Lower part)

230. Ferris clay pit, Ferris, Ellis County, Tex.
(P.P. 206)

70-T-1
(Ellis County)

Type locality for—

Spiroplectammina lalickeri Albritton & Phleger (J.P., vol.11,
p.353) 1937

Kyphopyxa cushmani Albritton & Phleger (J.P., vol.11, p.354) 1937

(P. lalickeri pl. 247)

Type locality for—
Spiroplectammina lalickeri Albritton & Phleger (J.P., vol. 11, p. 353)
Kyphopyxa cushmani Albritton & Phleger (J.P., vol. 11, p. 354)

Taylor group
(Lower part)

231. Clay pit at Palmer, Ellis County, Tex. J. A. Cushman and
J. A. Waters.

(P.P. 206)

70-T-2

(Ellis County)

Type locality for —

× Ventolabrella eggeri var. glabrata Cushman (C.C., vol.14, p.26, pl.4,
fig.19, 1938)

Type locality for —

Ventolabrella eggeri
var. glabrata Cushman (C.C. vol. 14, p. 26, '38)

Taylor group
(Lower part)

233. 2.3 miles by road north of Palmer, Ellis County, Tex.
J. A. Cushman and J. A. Waters.

(P.P. 206)

70-T-3

(Ellis County)

Type locality for--

15-4 '37 x Trochammia taylorana Cushman (C.C., vol.13, p.101)

Type locality for --
Trochammia taylorana (Cushman) (C.C., vol. 13, p. 101)

Taylor group
(Lower part)

234. Upper part. One and nine-tenths miles east of Bristol,
Ellis County, Tex., on road to State rock ferry on
Trinity River. C. H. Dane.

(P.P. 206)

70-T-4
(Ellis County)

Type locality for --

6-9, '33 x Dorothia glabella Cushman (C.C., vol.9, p.56)
16-9, '39 x Lituola irregulariter Cushman (C.C., vol.15, p.89)

Type locality for --
Dorothia glabella Cushman (C.C., vol.9, p.56)
Lituola irregulariter Cushman (C.C., vol.15, p.89)

Eagle Ford fm.

364. Shale about 175 feet below *Metoicoceras whitei* zone, 3 miles west of Midlothian, Ellis County, Tex. W. L. Moreman.

(P.F. 206)

70-T-5
(Ellis County)

Type locality for—

15-137 X Quinqueloculina moremani Cushman (C.C., vol 113, p.100)

Type locality for—
Quinqueloculina moremani Cushman
(C.C., vol. 13, p.100, '37)

Taylor fm.

70-T-6

(Ellis County)

Station 58. (Ellis County.)—Lower middle Taylor is exposed in a roadside ditch on the north side of the Garret-Boyce road, 2.8 miles by road east of the railroad station in the town of Boyce. This is the type locality for *Cythere taylorensis* n. sp.

p. 82, pl. 7, figs 3, 4 (U.T. Bull. 2907)

(Pummer Nov. 837)

Eagle Ford fm.

Station 39. (Ellis County.)—The *Metoicoceras whitei* zone of the Eagle Ford is exposed also in a northwestward-facing cliff on a tributary of Mountain Creek, one-quarter mile east of the concrete bridge on the Fort Worth-Midlothian highway about 3 miles north of Midlothian. Concentrates contain some sand, and whereas they are not rich in microfossils they contain a characteristic assortment of species.

(U.T. Bull. 2907)

Type locality for--

9-9,12

Cythereis eagelfordensis Alexander (U.T. Bull. 2907, p. 98, 1929)

70-1-7
(Ellis County)